

RAPIDLY DECAYING SUPERNOVA 2010X: A CANDIDATE “IA” EXPLOSION

MANSI M. KASLIWAL¹, S. R. KULKARNI¹, AVISHAY GAL-YAM², OFER YARON², ROBERT M. QUIMBY¹, ERAN O. OFEK¹, PETER NUGENT³, DOVI POZNANSKI^{3,12}, JANET JACOBSEN³, ASSAF STERNBERG², IAIR ARCAVI², D. ANDREW HOWELL^{4,16}, MARK SULLIVAN⁵, DOUGLAS J. RICH⁶, PAUL F. BURKE⁷, JOSEPH BRIMACOMBE MB ChB FRCA MD^{8,9}, DAN MILISAVLJEVIC¹⁰, ROBERT FESEN¹⁰, LARS BILDSTEN^{11,16}, KEN SHEN¹², S. BRADLEY CENKO¹², JOSHUA S. BLOOM¹², ERIC HSIAO³, NICHOLAS M. LAW¹³, NEIL GEHRELS¹⁴, STEFAN IMMLER¹⁴, RICHARD DEKANY¹⁵, GUSTAVO RAHMER¹⁵, DAVID HALE¹⁵, ROGER SMITH¹⁵, JEFF ZOLKOWER¹⁵, VISWA VELUR¹⁵, RICHARD WALTERS¹⁵, JOHN HENNING¹⁵, KAHNH BUI¹⁵ & DAN MCKENNA¹⁵

Draft version November 18, 2010

ABSTRACT

We present the discovery, photometric and spectroscopic follow-up observations of SN 2010X (PTF 10bhp). This supernova decays exponentially with $\tau_d = 5$ days, and rivals the current record-holder in speed, SN 2002bj. SN 2010X peaks at $M_r = -17$ mag and has mean velocities of 10,000 km s⁻¹. Our light curve modeling suggests a radioactivity powered event and an ejecta mass of 0.16 M_⊙. If powered by Nickel, we show that the Nickel mass must be very small (≈ 0.02 M_⊙) and that the supernova quickly becomes optically thin to γ -rays. Our spectral modeling suggests that SN 2010X and SN 2002bj have similar chemical compositions and that one of Aluminum or Helium is present. If Aluminum is present, we speculate that this may be an accretion induced collapse of an O-Ne-Mg white dwarf. If Helium is present, all observables of SN 2010X are consistent with being a thermonuclear Helium shell detonation on a white dwarf, a “Ia” explosion. With the 1-day dynamic-cadence experiment on the Palomar Transient Factory, we expect to annually discover a few such events.

Subject headings: supernovae: individual (SN 2010X, SN 2002bj), white dwarfs, surveys

1. INTRODUCTION

Our present knowledge of cosmic explosions is arguably biased by the searches themselves. In particular, the cadence and depth of many supernovae searches are designed to efficiently discover supernovae of type Ia (SNe Ia). A repeat visit to the sky on timescales of five days maximizes sky coverage and is still sufficient to catch SNe Ia on the rise. The brilliance of these events, peak absolute visual magnitude of -19 , sets the sensitivity of the searches. Conversely, fainter events and

those with a shorter characteristic lifetime are likely to be missed in such searches.

To illustrate the unexplored nature of this phase space, we plot the luminosity of optical transients versus their characteristic timescale (Figure 1). SNe Ia are confined to a narrow band (Phillips 1993) with decay timescales ranging from twelve days to three weeks. Classical novae span a large range of timescales albeit at considerably lower luminosities.

Figure 1 brings two white-spaces to attention: the wide “gap” in luminosity between novae and supernovae, and the apparent paucity of luminous events on short timescales.

Next, we discuss currently known exemplars of “faint” (i.e. lower luminosity than SNe Ia) and “fast” (i.e. faster than SN 2007ax) transients. SN 2005E occurred in the halo of its host galaxy and has been proposed as a Helium detonation on a binary white dwarf (Perets et al. 2010b). SN 2005cz has been proposed to have a massive star origin (Kawabata et al. 2010). SN 2008ha is also being widely debated both as a deflagration of a white dwarf (Foley et al. 2009b,a) and core-collapse of a massive star (Valenti et al. 2009).

Until recently, the fastest event known was SN 2002bj (Poznanski et al. 2010). It decayed by one magnitude in five days and was quite spectroscopically peculiar. The origin of this event is not yet clear.

The Palomar Transient Factory¹⁷ (PTF) was motivated in great measure to systematically explore the phase space for fast and faint explosive transients (Law et al. 2009; Rau et al. 2009). Here, we present the discovery of a fast event, SN 2010X (PTF 10bhp).

¹ Cahill Center for Astrophysics, California Institute of Technology, Pasadena, CA, 91125, USA

² Ben-Zvi Center for Astrophysics, Faculty of Physics, The Weizmann Institute of Science, Rehovot 76100, Israel

³ Computational Cosmology Center, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA

⁴ Las Cumbres Observatory Global Telescope Network, Inc, Santa Barbara, CA, 93117, USA

⁵ Department of Physics, Oxford University, Oxford, OX1 3RH, UK

⁶ Rich Observatory, 62 Wessnette Dr., Hampden, Maine USA

⁷ Burke Observatory, 19 Berry Rd., Pittsfield, Maine USA

⁸ New Mexico Skies Observatory, Mayhill New Mexico, USA

⁹ James Cook University, Cairns, Australia

¹⁰ 6127 Wilder Lab, Department of Physics and Astronomy, Dartmouth College, Hanover, NH, 03755, USA

¹¹ Kavli Institute of Theoretical Physics, University of California Santa Barbara, Santa Barbara, CA 93106, USA

¹² Department of Astronomy, 601 Campbell Hall, University of California, Berkeley, CA 94720-3411, USA

¹³ Dunlap Institute for Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto M5S 3H4, Ontario, Canada

¹⁴ NASA-Goddard Space Flight Center, Greenbelt, MD 20771, USA

¹⁵ Caltech Optical Observatories, California Institute of Technology, Pasadena, CA 91125, USA

¹⁶ Department of Physics, University of California Santa Barbara, Santa Barbara, CA 93106, USA

¹⁷ <http://www.astro.caltech.edu/ptf>




FIG. 1.— **Left:** The phase space of cosmic explosive transients. The color for each event represents the color at peak brightness. The band to the top right denotes supernovae of type Ia. The fastest such event is SN 2007ax (Kasliwal et al. 2008). Classical novae occupy a band between -6 and -10 magnitude. Note that the only two transients with a timescale shorter than ten days are SN 2010X (PTF 10bhp) and SN 2002bj. **Right:** The multi-band optical light curve of SN 2010X (colored circles; green is g-band, red is r-band, orange is i-band). Three white light measurements have been calibrated to r-band and denoted by red circles with blue outline. Downward arrows represent upper limits. All light curves are normalized and shifted so that peak magnitude is zero and the time at peak is set to zero. For SN 2010X the epoch of maximum light is at MJD of 55239.5. The fast evolution of SN 2010X is compared to the current recordholder for fast supernovae, SN2002bj (gray triangles; r-band; Poznanski et al. 2010). Also shown is a prototypical “fast” Type Ic supernova, SN1994I (dashed line; Richmond et al. 1996) and templates^a of the fast Type Ia SN1991bg and slow Type Ia SN1991T (Nugent et al. 2002). Note the rapid rise and the spectacular decay of SN 2010X and SN 2002bj relative to the other Type I exemplars.

^ahttp://supernova.lbl.gov/~nugent/nugent_templates.html

2. DISCOVERY

On UT 2010 February 7.07, D. Rich of Hampden, Maine discovered a transient in the galaxy NGC 1573A at RA(J2000)=04^h48^m27.7^s and Dec(J2000)=+73°28′13″. The discovery was confirmed by P. Burke of Pittsfield, Maine, upon which a notification was issued (CBET 2166; Rich & Burke 2010) and the transient dubbed SN2010X. On UT 2010 February 19.13, the Palomar Transient Factory independently detected this same transient and the pipeline assigned the name, PTF 10bhp. PTF had previously undertaken observations of this field (as a part of the dynamic cadence experiment) on January 11, 17 and 25 but with no detection.

3. OPTICAL LIGHT CURVE

Energized by the apparent rapid fading, we initiated follow-up observations. The photometric observations from the 2-m Faulkes North Telescope (FTN) of the Las Cumbres Observatory Global Telescope (LCOGT), PTF, the Palomar Hale 200-inch telescope (P200) as well as white light observations provided by our amateur astronomer colleagues are summarized in Figure 1.

SN 2010X is located close to the nucleus of its host galaxy (4.4″ E, 6.0″ N) and as such galaxy light subtraction is critical to produce reliable photometry. The images were subtracted from a template image using the software `hotpants` and `wcsremap` to measure a convolution kernel and align the images respectively (both codes

supplied by A. Becker¹⁸). Aperture photometry was performed on each of these in a self-consistent manner using the same set of 22 calibration stars. Conversions from USNO-B1 magnitudes to SDSS gri magnitudes were done adopting Jordi et al. (2006). The resulting light curve is plotted in Figure 1.

Overplotting SN 2002bj, we find that light curves of the two supernovae are remarkably similar. Linearly fitting all the r-band detections post maximum light, we measure that SN 2010X decayed by 0.23 ± 0.01 mag day⁻¹. The corresponding exponential timescale (in the r-band) is $\tau_d = 4.7 \pm 0.2$ days.

The foreground Galactic extinction along the line of sight is $E(B-V)=0.146$ or $A_r=0.4$ (Schlegel et al. 1998). The redshift of NGC 1573A is 0.015. Assuming standard cosmology (and $h_0=0.72$), we adopt a distance of 62.5 Mpc and a distance modulus of 34.0. Thus, the peak absolute magnitude of SN 2010X is $M_r \approx -17.0$ mag, 1.5 mag less luminous than SN2002bj.

4. SPECTROSCOPIC FOLLOW-UP

On February 8 and 9, the first spectra (Figure 2) to classify the nature of this transient were taken with CCDs on the 2.4m Hiltner telescope of the MDM observatory (CBET 2167, Milisavljevic & Fesen 2010). Comparison with a library of supernova spectra using SNID (Blondin & Tonry 2007) showed resemblance to the Type Ic supernovae SN 1994I and SN 2004aw a few days before maximum light. Further observations (Fig-

¹⁸ http://www.astro.washington.edu/users/becker/c_software.html




FIG. 2.— Spectroscopic follow-up of SN 2010X by MDM, Gemini, Keck and Palomar Observatories. The phase of the spectra relative to maximum light is labeled. Note the velocity evolution. Also shown is a spectrum of SN 2002bj (Poznanski et al. 2010).

ure 2) were undertaken on Gemini-North/GMOS (Feb 23), Keck I/LRIS (Mar 7) and the Hale 200-inch/DBSP (Mar 18) telescopes. No perfect matches to Ic (or Ia, Ib) templates were found for these spectra. The velocity evolved from 12000 km s^{-1} before maximum to 9000 km s^{-1} at late-time.

We used SYNOW (Jeffery & Branch 1990) to infer elements in the spectra of SN 2010X (Figure 3). The most prominent identifications are oxygen (O I lines), Calcium (both Ca II IR triplet and Ca II H+K on the blue side), Carbon (C II lines), Titanium (Ti II) and Chromium (Cr II). Ti II and Cr II explain the broad blue features and adding Fe II improves the fit slightly. There is also some evidence for Mg I albeit based on single line.

The presence of Helium (He I), Sodium (Na D) and Aluminum (Al II) is less clear and we illustrate this dilemma in the inset of Figure 3. He I has three relevant lines: 5876 Å , 6678 Å and 7065 Å . The absorption feature around 5700 Å can be explained by both He I as well as Na D. The absorption feature around 6850 Å can be explained by Al II or He I. Since the central He I line is not prominent, SYNOW suggests that the combination of Na D and Al II is a better fit. However, Branch (2003) discuss that this central He I line is a singlet transition and this may both be suppressed and blueshifted in non-LTE relative to the other two He I triplet transitions. Therefore, we cannot conclusively say whether or not Helium is present in SN 2010X.

Comparing the spectra, SN 2002bj has substantially lower velocities (4000 km s^{-1} at +7d vs. $10,000 \text{ km s}^{-1}$ at +10d) and a bluer continuum ($g-r=0.2$ at +12d vs. $g-r=1.2$ at +23d) than SN 2010X. Consistent with the SYNOW fit shown in Poznanski et al. 2010, the elements in common between the two supernovae are O I, C II and




FIG. 3.— SYNOW fit to the Keck spectrum (+23 days) of SN 2010X. Lines contributed by each ion are shown. Fits with (purple) and without (red) Iron are overplotted on the data (blue). **Top Panel:** The dilemma of whether SN 2010X has Helium or a combination of Sodium and Aluminum. The vertical dashed lines show Helium at 9500 km s^{-1} . In non-LTE, the singlet transition of $\lambda 6678$ may be suppressed relative to the $\lambda 7065$ and $\lambda 5876$ triplet transition Helium lines.

Mg II. The primary difference is the presence of Ca II in SN 2010X and presence of S II in SN 2002bj. We re-fit the spectrum of SN 2002bj with the same elements as in SN 2010X. We find that the presence of Al II vs. He I is just as ambiguous for SN 2002bj as SN 2010X. Similar to SN 2010X, including Fe II improves the fit but the presence of Fe-group elements in SN 2002bj is not conclusive.

5. MODELING THE LIGHT CURVE

The excellent match between the normalized light curves of SN 2010X and SN 2002bj (see Figure 1) suggests that these two SNe belong to the same class of explosions. Combining the two data sets allows a robust determination of the rise time¹⁹ ($\tau_r \approx 6 \text{ d}$) and the subsequent exponential decay ($\tau_d \approx 5 \text{ d}$)

The peak bolometric luminosity of SN 2010X is $L_{\text{peak}} = 10^{42} \text{ erg s}^{-1}$. While the expansion speed varies from $12,000 \text{ km s}^{-1}$ at early times to $9,000 \text{ km s}^{-1}$ at late times, we accept $v_s \approx 10,000 \text{ km s}^{-1}$ as a representative value.

The rise time in an explosion is the geometric mean of the initial photon diffusion timescale and the initial hydrodynamic time scale²⁰. Thus, $\tau_r^2 \propto \kappa M_{\text{ej}} / v_s$ where κ is the opacity. Assuming that the mean opacity of SN 2010X is the same as that for SNeIa events, (for which, following Hayden et al. 2010, we adopt the following: $M_{\text{ej}} \approx 1.4 M_{\odot}$, $v_s = 10^9 \text{ cm s}^{-1}$ and $\tau_r \approx 17.5 \text{ d}$), we obtain $M_{\text{ej}} \approx 0.16 M_{\odot}$. This gives an explosion en-

¹⁹ time from explosion to peak brightness

²⁰ The derivation can be found in the textbooks, Arnett 1996 and Padmanabhan 2001.

ergy, $E_0 = 1/2 M_{\text{ej}} v_s^2 \approx 1.7 \times 10^{50}$ erg.

Next, we investigate a physical model that satisfactorily accounts for the rise time, the decay time, the peak luminosity and the expansion velocity.

5.1. Pure explosion

The simplest model is an explosion in which all the explosive energy (E_0) is deposited instantaneously into the ejecta. The peak luminosity is then $E_0/t_d(0)$ where $t_d(0) \propto \kappa M_{\text{ej}}/R_0$ is the initial photon diffusion time; here, R_0 is the radius of the progenitor. Following peak luminosity, the decay is rapid: $\log(L) \propto -(t/\tau_r)^2$. The virtue of this model is that one can obtain an arbitrarily rapid rate of decay since, over any limited stretch of time, the light curve can be approximated by a linear decay with the desired value for the slope.

For SN 2010X, we find $R_0 \sim 4 \times 10^{12}$ cm. The large inferred radius would make sense if the progenitor had an envelope (as in type II supernovae). The absence of hydrogen at any phase of the supernova (see §4) argues strongly against this model. Hence, we reject this hypothesis.

5.2. Radioactivity powered explosion

The next level of models is that developed for SNe Ia explosions, where the peak luminosity and subsequent decay is governed by radioactive material present in the ejecta. In this model, expansion decreases the store of internal energy whereas radioactivity increases it. If the photon diffusion time-scale is long, most of the radioactive energy goes into expansion. Once the diffusion time-scale becomes smaller than the expansion time-scale, the light curve tracks the radioactive luminosity (Arnett 1982), provided that there is sufficient optical depth for the γ -rays emitted during radioactive decay to undergo multiple scatterings and lose their energy to electrons.

The primary source of luminosity in a SNe Ia model is the heat provided by γ -rays emitted as ^{56}Ni decays to ^{56}Co and then to ^{56}Fe . In SNe Ia, the column density of the ejecta is thick enough to trap the γ -rays and successive Compton scatterings extract energy from the γ -rays (at least for the first month). However, given the small ejecta mass for SN 2010X, attention has to be paid to the possibility that γ -rays from decaying nuclei may escape without depositing their energy into the ejecta.

The electron (Thompson) optical depth is:

$$\tau_e = n_e R \sigma_T = \frac{3}{4\pi} \frac{M_{\text{ej}}}{m_p} \frac{Z}{A} \frac{\sigma_T}{R^2} \sim 9 \left(\frac{M_{\text{ej}}}{0.16 M_\odot} \right) \left(\frac{t}{15 \text{ day}} \right)^{-2} \quad (1)$$

where Z is the atomic number, A is the mass number, m_p is mass of proton, σ_T is the Thompson cross-section and $R \sim 6(t/\text{day}) \text{ AU}$ is the radius at time t .

Thus, there appears to be sufficient optical depth at the epoch of peak luminosity to trap most of the γ -rays. Thus, for SN 2010X, the peak luminosity of $10^{42} \text{ erg s}^{-1}$ corresponds to ^{56}Ni mass of about $0.02 M_\odot$ – a very small amount by the standards of most supernovae. For SN 2002bj, the peak luminosity was $10^{43} \text{ erg s}^{-1}$ (Poznanski et al. 2010) and the inferred ^{56}Ni mass was correspondingly larger, $0.2 M_\odot$.

Next, we use a fitting formula (as given in Kulkarni 2005; Equation 47) to estimate the fraction of γ -rays which are effectively absorbed inside the ejecta, $\eta(\tau_e)$. The kinetic energy of positrons (3.5% of L_{Co} ; Sollerman et al. 2002) dominates by day 51. Hence, the radiated luminosity, $L_{\text{rad}} = (0.965\eta + 0.035)L_{\text{Co}} + \eta L_{\text{Ni}}$ where L_{Ni} is the radioactive power released by the decay of ^{56}Ni , and L_{Co} by the daughter ^{56}Co . In Figure 4, we display the luminosity due to radioactivity and that actually trapped in the ejecta — the latter shows a satisfactory agreement with the observations.




FIG. 4.— Shown above is the radioactive luminosity (solid line) and absorbed luminosity (dashed line) for the following model parameters: $M_{\text{ej}} = 0.16 M_\odot$, $M_{\text{Ni}} = 0.02 M_\odot$ and $v = 10^9 \text{ cm s}^{-1}$. Also shown is a quasi-bolometric light curve of SN 2010X estimated by (a) computing νF_ν in r -band (empty circles are detections, inverted triangles are upper limits), and (b) integrating the optical spectrum (filled circles). Also shown is a comparison to a “Ia” light curve (red dotted line; Shen et al. 2010) assuming: $M_{\text{wd}} = 1.2 M_\odot$, $M_{\text{env}} = 0.05 M_\odot$, $M_{\text{ej}} = 0.036 M_\odot$, $M_{\text{Fe}} = 0.005 M_\odot$, $M_{\text{Ni}} = 0.02 M_\odot$, $M_{\text{Cr}} = 0.0002 M_\odot$.

5.2.1. Possible X-ray signature

An optically thin ejecta opens up the possibility of detecting the γ -rays (or degraded hard X-rays) emitted during β -decay. The *Swift* Observatory observed SN 2010X for 9758.7 s on MJD 55248.775 (9 days past peak). We constrain the X-ray flux²¹ to be less than $0.00050 \text{ counts s}^{-1}$ or $7.7 \times 10^{39} \text{ erg s}^{-1}$. By this epoch, our model shows that $L_\gamma \sim 10^{41} \text{ erg s}^{-1}$. Since photon number is conserved in scattering, the luminosity in the *Swift* band is expected to be a factor of 200 smaller and hence, the upper limit is not constraining.

6. ENVIRONMENT

The host of SN 2010X, NGC 1573A, is a small ($1.6'$ diameter), spiral galaxy variously classified as Sb (UGC) and SABbc (RC3). The host of SN 2002bj, NGC 1821, is a small ($1.1'$ diameter), barred irregular galaxy classified as IB(s)m. Both transients occurred close to the

²¹ we note that the six photons in the XRT HPD PSF of $0.3'$ are likely from the galaxy nucleus

galaxy nucleus — 2.3 kpc for SN 2010X and 1.8 kpc for SN 2002bj. In Figure 5, we show the location of the supernovae in deep images of the galaxy.




FIG. 5.— *Left:* R-band image of NGC 1573A, the host of SN 2010X, taken with the Large Format Camera on the Palomar 200-inch telescope. *Right:* Sum of all available Deepsky^a (Nugent 2009) images of NGC 1821, the host of SN 2002bj.

^a<http://supernova.lbl.gov/~nugent/deepsky.html>.

7. CONCLUSION

To summarize, SN 2010X is the second member of a class of supernovae that declines exponentially on timescales shorter than 5 days. Relative to SN 2002bj, SN 2010X is less luminous by 1.5 mag ($M_R \approx -17$) and has higher velocities ($10,000 \text{ km s}^{-1}$) by more than a factor of two. Both events have a small inferred ejecta mass. Both events are spectroscopically different from any other type I supernovae. The spectra for both supernovae can be modeled with mostly similar elements (C, O, Mg, Si, Ti and Fe). The evidence (or lack thereof) for Helium is not conclusive in both cases.

If SN 2010X is powered by radioactive ^{56}Ni , the combination of a rapid rise time and low peak luminosity constrains the Nickel mass to be small, $0.02 M_\odot$. ^{56}Ni constitutes $\approx 13\%$ of ejecta. However, under the same assumptions, ^{56}Ni would constitute bulk of the ejecta mass for SN 2002bj. Thus, while in both cases the ejecta mass remains the same, the nucleo-synthesis may be strongly variable. We also show that given the small ejecta mass, γ -rays from decaying ^{56}Ni can start escaping from the ejecta shortly after peak brightness. This early escape reasonably accounts for the rapid decay of the light curve of SN 2010X.

Perets et al. (2010a) have argued that S Andromeda (the first recorded SN in Andromeda) and SN 1939B (the first recorded SN in Virgo) are also like SN 2002bj. The claim primarily rests on rapid rise and rapid decay at early-time. It is of some interest to note that the late-time (2 months to nearly 1 year) decay rates, 0.03 mag/day for S Andromeda and 0.02 mag/day for SN 1939B (Perets et al. 2010a), are consistent with ^{56}Co decay (with some escape of γ -rays). A consistent explanation would require a two-zone model: comparable amount of ^{56}Ni in a slowly expanding core (to account for the late time light curve) and a rapidly expanding shell (to account for the rapid decay seen after peak brightness).

An alternative model is that the early-time emission is powered by another suitably rapidly decaying radioactive element(s). If powered solely by ^{48}Cr , $0.02 M_\odot$

is adequate. Recently, Shen et al. 2010 computed models and observables for “Ia” explosions (Bildsten et al. 2007) powered by ^{48}Cr , ^{52}Fe and ^{56}Ni : rise time between 2–10 days, ejecta velocity between $9000\text{--}13000 \text{ km s}^{-1}$, peak luminosity between $0.5\text{--}5 \times 10^{42} \text{ erg s}^{-1}$ and presence of Ca II and Ti II in the spectra. The properties of SN 2010X are consistent with all these predictions. Specifically, the light curve model presented for a core mass of $1.2 M_\odot$ and envelope mass of $0.05 M_\odot$ is a reasonable match (Figure 4). Furthermore, Shen et al. 2010 also discuss that the presence of Helium in the spectra may be a non-LTE effect.

If Aluminum is indeed present in the spectra, the avenue for a speculative scenario opens up. Neither ^{26}Al nor ^{27}Al is a product of Helium burning. Aluminum can be made via explosive burning of Neon and/or Carbon (Arnett & Bazan 1997; Woosley & Weaver 1980). Perhaps, SN 2010X is the outcome of accretion induced collapse of an O-Ne-Mg white dwarf (Metzger et al. 2009).

Finally, we note that the rich, star-forming environment of SN 2010X and SN 2002bj does not preclude a massive star origin. Fallback events, where a massive star collapses into a black hole, are also expected to be fast declining (Fryer et al. 2009; Moriya et al. 2010). However, the velocities expected from these models are significantly lower than observed and the spectra are more substantially dominated by intermediate mass elements.

Regardless of all these rich possibilities, it is clear that further progress in understanding the nature of these ephemeral transients would require a larger sample. Fortunately, PTF, especially as it moves to “dynamic” 1-day cadence (Law et al. 2009) targetting nearby galaxies and clusters, is well equipped to annually find a few such events. Late-time photometry is important to look for tell-tale signatures of ^{56}Co decay. Sensitive optical (or better still, ultra-violet) spectroscopy may directly reveal the radioactive element(s) powering these events. It is also not inconceivable (given the history of S Andromeda and SN 1939B) that we will be lucky enough to observe a local analog of such an event with the hard X-ray mission, NuSTAR (Harrison et al. 2010).

Acknowledgments. MMK thanks the Gordon and Betty Moore Foundation for a Hale Fellowship in support of graduate study. MMK thanks the Purnarth Headquarters in Indore, India for their warm hospitality while writing this manuscript.

We thank the referee for accepting this paper on a timescale faster than that characterizing the rapid photometric evolution of this supernova!

We would like to acknowledge the following discussions: MMK & Paolo Mazzali, AGY & David Branch, SRK & Xiaofeng Wang, DAH & Ryan Foley, MMK & Rollin Thomas. We are grateful to the staff of the Gemini Observatory and Swift Observatory for efficiently executing TOO triggers. We thank the librarians who maintain the ADS, the NED, and SIMBAD data systems.

The Weizmann Institute PTF participation is supported by grants to AGY from the Israel Science Foundation and the US-Israel Binational Science Foundation. EOO and DP are supported by an Einstein Fellowship. SBC is grateful for support from Gary and Cynthia Bengier and the Richard and Rhoda Goldman Fund. Computational resources and data storage were

contributed by NERSC, supported by U.S. DoE contract DE-AC02-05CH11231. PEN acknowledges support from the US DoE contract DE-FG02-06ER06-04.

Facilities: PO:1.2m (), Hale (), Gemini:Gillett (),Hiltner (),Swift ()

REFERENCES

- Arnett, D. 1996, Supernovae and nucleosynthesis. an investigation of the history of matter, from the Big Bang to the present, ed. Arnett, D.
- Arnett, D., & Bazan, G. 1997, *Science*, 276, 1359
- Arnett, W. D. 1982, *ApJ*, 253, 785
- Bildsten, L., Shen, K. J., Weinberg, N. N., & Nelemans, G. 2007, *ApJ*, 662, L95
- Blondin, S., & Tonry, J. L. 2007, *ApJ*, 666, 1024
- Branch, D. 2003, in *IAU Symposium*, Vol. 212, A Massive Star Odyssey: From Main Sequence to Supernova, ed. K. van der Hucht, A. Herrero, & C. Esteban, 346–+
- Foley, R. J., Brown, P. J., Rest, A., Challis, P. J., Kirshner, R. P., & Wood-Vasey, W. M. 2009a, *ArXiv e-prints*
- Foley, R. J., et al. 2009b, *AJ*, 138, 376
- Fryer, C. L., et al. 2009, *ApJ*, 707, 193
- Harrison, F. A., et al. 2010, *ArXiv e-prints*
- Hayden, B. T., et al. 2010, *ApJ*, 712, 350
- Jeffery, D. J., & Branch, D. 1990, in *Supernovae*, Jerusalem Winter School for Theoretical Physics, ed. J. C. Wheeler, T. Piran, & S. Weinberg, 149
- Jordi, K., Grebel, E. K., & Ammon, K. 2006, *A&A*, 460, 339
- Kasliwal, M. M., et al. 2008, *ApJ*, 683, L29
- Kawabata, K. S., et al. 2010, *Nature*, 465, 326
- Kulkarni, S. R. 2005, *ArXiv Astrophysics e-prints*
- Law, N. M., et al. 2009, *PASP*, 121, 1395
- Metzger, B. D., Piro, A. L., Quataert, E., & Thompson, T. A. 2009, *ArXiv e-prints*
- Milisavljevic, D., & Fesen, R. 2010, *Central Bureau Electronic Telegrams*, 2167, 1
- Moriya, T., Tominaga, N., Tanaka, M., Nomoto, K., Sauer, D. N., Mazzali, P. A., Maeda, K., & Suzuki, T. 2010, *ApJ*, 719, 1445
- Nugent, P., Kim, A., & Perlmutter, S. 2002, *PASP*, 114, 803
- Nugent, P. E. 2009, in *Bulletin of the American Astronomical Society*, Vol. 41, *Bulletin of the American Astronomical Society*, 419
- Padmanabhan, T. 2001, *Theoretical Astrophysics, Volume 2: Stars and Stellar Systems*, ed. Padmanabhan, T.
- Perets, H. B., Badenes, C., Arcavi, I., Simon, J. D., & Gal-yam, A. 2010a, *ArXiv e-prints*
- Perets, H. B., et al. 2010b, *Nature*, 465, 322
- Phillips, M. M. 1993, *ApJ*, 413, L105
- Poznanski, D., et al. 2010, *Science*, 327, 58
- Rau, A., et al. 2009, *PASP*, 121, 1334
- Rich, D., & Burke, P. 2010, *Central Bureau Electronic Telegrams*, 2166, 1
- Richmond, M. W., et al. 1996, *AJ*, 111, 327
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Shen, K. J., Kasen, D., Weinberg, N. N., Bildsten, L., & Scannapieco, E. 2010, *ApJ*, 715, 767
- Sollerman, J., et al. 2002, *A&A*, 386, 944
- Valenti, S., et al. 2009, *Nature*, 459, 674
- Woosley, S. E., & Weaver, T. A. 1980, *ApJ*, 238, 1017

